

Retinoic acid generates a beneficial microenvironment for liver progenitor cell activation in acute liver failure.

Info

PUBLIZIERT AM JOURNAL
2024-08-01 Hepatol Commun

LINK

<https://www.ncbi.nlm.nih.gov/pubmed/39023343>

Autoren

S. Wang F. Link S. Munker W. Wang R. Feng
R. Liebe Y. Li Y. Yao H. Liu C. Shao
M. P. A. Ebert H. Ding S. Dooley H. L. Weng
S. S. Wang

Abstract

BACKGROUND: When massive necrosis occurs in acute liver failure (ALF), rapid expansion of HSCs called liver progenitor cells (LPCs) in a process called ductular reaction is required for survival. The underlying mechanisms governing this process are not entirely known to date. In ALF, high levels of retinoic acid (RA), a molecule known for its pleiotropic roles in embryonic development, are secreted by activated HSCs. We hypothesized that RA plays a key role in ductular reaction during ALF. **METHODS:** RNAseq was performed to identify molecular signaling pathways affected by all-trans retinoid acid (atRA) treatment in HepaRG LPCs. Functional assays were performed in HepaRG cells treated with atRA or cocultured with LX-2 cells and in the liver tissue of patients suffering from ALF. **RESULTS:** Under ALF conditions, activated HSCs secreted RA, inducing RARalpha nuclear translocation in LPCs. RNAseq data and investigations in HepaRG cells revealed that atRA treatment activated the WNT-beta-Catenin pathway, enhanced stemness genes (SOX9, AFP, and others), increased energy storage, and elevated the expression of ATP-binding cassette transporters in a RARalpha nuclear translocation-dependent manner. Further, atRA treatment-induced pathways were confirmed in a coculture system of HepaRG with LX-2 cells. Patients suffering from ALF who displayed RARalpha nuclear translocation in the LPCs had significantly better MELD scores than those without. **CONCLUSIONS:** During ALF, RA secreted by activated HSCs promotes LPC activation, a prerequisite for subsequent LPC-mediated liver regeneration.