

Transforming Growth Factor- β Signaling in Alcohol-Associated Liver Disease

Info

PUBLIZIERT AM

2025-10-01

JOURNAL

The American Journal of Pathology

LINK

<https://doi.org/10.1016/j.ajpath.2025.09.017>

Autoren

Huihui Zou Sai Wang Chenjun Huang

Steven Dooley Nadja M. Meindl-Beinker

Abstract

Transforming growth factor- β (TGF- β) signaling exerts broad regulatory effects on alcohol-associated liver disease (ALD) progression, influencing processes such as hepatocellular injury, regeneration, inflammation, fibrogenesis, cirrhosis, carcinogenesis, and hepatic failure. TGF- β modifies alcohol-induced signals in multiple liver-resident cell types, including hepatocytes, hepatic stellate cells, liver sinusoidal endothelial cells, and immune populations, particularly macrophages. To delineate its context-specific roles in ALD, 154 of 421 PubMed-listed publications (2000 to 2025; search terms TGF- β and alcohol and liver disease) were reviewed, supplemented by 19 foundational studies published earlier. In hepatocytes, TGF- β promotes oxidative stress, apoptosis, metabolic reprogramming, and epithelial-to-mesenchymal transition. In hepatic stellate cells and Kupffer cells, gut-derived endotoxins, ethanol, and unsaturated fatty acids induce TGF- β alongside proinflammatory cytokines. Ethanol metabolism generates acetaldehyde, which drives TGF- β and receptor expression, enhances canonical and noncanonical signaling, and engages epigenetic regulators to promote extracellular matrix deposition. In liver sinusoidal endothelial cells, alcohol-induced TGF- β suppresses proliferation, contributing to sinusoidal capillarization, impaired endothelial regeneration, and fibrogenesis. TGF- β dampens clearance of damaged hepatocytes and perpetuating chronic injury by suppressing natural killer cell cytotoxicity and promoting regulatory T-cell differentiation. At end-stage disease, TGF- β promotes expansion and fate switching of cholangiocyte-derived liver progenitor cells to replenish lost hepatocytes. Despite its central role in ALD, therapeutic exploitation of TGF- β signaling remains underexplored. Future studies should define cell type-specific signaling nodes to enable precision therapies.