NADPH oxidase 1/4 dual inhibition impairs transforming growth factor-beta protumorigenic effects in cholangiocarcinoma cancer-associated fibroblasts

Info

PUBLIZIERT AM
2025-12-01

JOURNAL
Signal Transduction and Targeted Therapy

LINK
https://doi.org/10.1038/s41392-025-02347-z

Autoren

Josep Amengual Ester Gonzalez-Sanchez Mariana Yáñez-Bartolome Laura Sererols-Viñas Aashreya Ravichandra Celia Guiton Noel P. Fuste Ania Alay Sara Hijazo-Pechero Beatriz Martín-Mur Marta Gut Anna Esteve-Codina Ana Cantos-Cortes Rut Espinosa-Sotelo Emilio Ramos Teresa Serrano Mariona Calvo Berta Laquente Joana Ferrer Gabriel Pons Andrés Mendez-Lucas Steven Dooley Sumera I. Ilyas Marie Vallette Lynda Aoudjehane Marie Lequoy Laura Fouassier Cédric Coulouarn Silvia Affò Alexander Scheiter Diego F. Calvisi Tian V. Tian Isabel Fabregat Javier Vaquero

Abstract

Transforming growth factor beta (TGF-β) signalling has become an attractive therapeutic target due to its pro-tumorigenic actions on epithelial cells and its immunosuppressive effects in the tumour microenvironment. In intrahepatic cholangiocarcinoma (iCCA), a highly aggressive malignancy of the biliary tract with poor prognosis, the latest clinical trials using TGF-β inhibitors have failed indicating that the specific actions carried out by TGF-β in iCCA are yet not well delineated. Here, we show that TGF-β signalling is highly active in iCCA and exerts a prominent suppressor effect on tumour cell lines and organoids established from iCCA metastases biopsies, that relies on a functional canonical SMAD2/3/4 signalling. Thus, TGF-β inhibitors promote, instead of inhibiting, tumour cell growth. In this context, a promising strategy is to target intracellular proteins downstream the TGF-β receptors accounting only for TGF-β pro-tumorigenic actions. NADPH oxidase 4 (NOX4), a downstream mediator of the TGF-β signalling pathway, is strictly expressed in cancer-associated fibroblasts (CAF) of iCCA and acts in concert with NOX1 to regulate CAF functions. Use of a dual NOX4/NOX1 inhibitor impaired CAF actions and reduced tumour growth in vitro and in two different in vivo iCCA experimental models. Collectively, our findings reveal an actionable way to specifically target TGF-β pro-tumorigenic actions in CAF from iCCA without undesirable side effects on tumour cells, suggesting a potentially bright future for dual NOX4/NOX1 inhibitors in the clinics, alone or in combination with other therapies.