Comparison and evaluation of methods to infer gene regulatory networks from multimodal single-cell data

Info


PUBLIZIERT AM
2024-12-21

JOURNAL
preprint

LINK
https://doi.org/10.1101/2024.12.20.629764

Autoren

Pau Badia-i-Mompel Roger Casals-Franch Lorna Wessels Sophia Müller-Dott Rémi Trimbour Yunxiao Yang Ricardo O. Ramirez Flores Julio Saez-Rodriguez



Abstract

Cells regulate their functions through gene expression, driven by a complex interplay of transcription factors and other regulatory mechanisms that together can be modeled as gene regulatory networks (GRNs). The emergence of single-cell multi-omics technologies has driven the development of several methods that integrate transcriptomics and chromatin accessibility data to infer GRNs. While these methods provide examples of their utility in discovering new regulatory interactions, a comprehensive benchmark evaluating their mechanistic and predictive properties as well as their ability to recover known interactions is lacking. To address this, we built a comprehensive framework, Gene Regulatory nETwork Analysis (GRETA), available as a Snakemake pipeline, that includes state of the art methods decomposing their different steps in a modular manner. With it, we found that the GRNs were highly sensitive to methods' choices, such as changes in random seeds, or replacing steps in the inference pipelines, as well as whether they use paired or unpaired

multimodal data. Although the obtained networks performed well in predictive evaluation tasks and partially recovered known interactions, they struggled to capture causal relationships from perturbation assays. Our work brings attention to the challenges of inferring GRNs from single-cell omics, offers guidelines, and presents a flexible framework for developing and testing new approaches.